Diagram 2:


COLEVA
Known and Suspected Consequences Of Lifetime Exposures to Violence and Abuse
Consequences of Lifetime Abuse
Ophthalmology
Traumatic injuries
Transient blindness
Double vision
Endocrine
Elevated T3/levels
Oxytocin-vasopressin
ACTH-Glucocorticoids
Prolactin
Infectious disease
Increased risk of HIV
Hepatitis B and C
STIs
ENT
Dizziness
Ruptured TM
Nasal fractures
Mandible fractures
TMJ disorders
General
Chronic fatigue
Eating disorders
Obesity
Anorexia
Bulimia
Sleep disorders
Insomnia
Hypersomnia
Risk of experiencing violence and abuse
Risk of perpetrating violence and abuse
Cardiovascular
Palpitations
Atypical chest pains
Increased risk of CAD
Gastrointestinal
Gastritis
Unexplained vomiting
Pancreatitis
Liver disease
Irritable bowel syndrome
Chronic abdominal pain
GERD
Encopresis
Ob-Gyn
Chronic pelvic pain
Increased STIs
Teen Pregnancies
Abortion
Spontaneous
Therapeutic
Infertility
Premenstrual dysphoric disorder
Pelvic inflammatory disease
Preterm labor and low birth weight
Hyperemesis gravidarum
Complications of labor and delivery
Prolonged labor
Greater need for medication
Increased risk of C-section
Post-partum depression
Anorgasmia
Sexual dysfunction
Hypersexual
Unnecessary surgical procedures
Less frequent self-breast exam
Increase risk of cervical cancer
Dyspareunia
Vulvodynia
Vaginismus
Altered ovarian function
Early menopause
Allergies
Multiple medication intolerance
Environmental sensitivities
Surgery
Increased surgical procedures in general
More complications from surgery
Higher rate of surgical failure
Orthopedics
Chronic low back pain
Increased risk of any fracture
Neck strain - radiculopathy
Complex regional pain
Rheumatology
Fibromyalgia
Chronic fatigue
Auto-immune disorders
Rheum. Arthritis
Neurological
Chronic Daily Headaches
Seizures
Pseudoseizures
Syncope
Altered sensations
Transient amnesia
Increased risk of CVA
Primary CVA-strangle
Secondary to high risk behavior
Dizziness/ lightheadedness
GU
Urethritis
Bladder dysfunction
Prostatitis
Epididymitis
Frequent bladder infections
Interstitial cystitis
Dental
TMJ Disorders
Bruxism
Poor dental hygiene
Increased caries
Gingivitis
Dental abscesses
Increased tooth extraction rate
Avoidance of dental care
Dermatology
Unexplained pruritis
Chronic or recurring urticaria
Chronic excoriation - picking
Alopecia - trichotillomania
Neurodermatitis
Delusional Parasitosis
Mental Health
Depression
Anxiety
Panic disorder
PTSD
Bipolar disorder
Cutting and other self-inflicted injury
Suicidal/suicidal attempts
Aggression/anger problems
Agoraphobia
Alexithymia
Munchausen's syndrome
Somatization syndrome
Dissociative disorders
Obsessive-compulsive disorder
Hallucinations
Schizophrenia
Behavioral Issues
Alcoholism
Illicit drug use
Tobacco use
Decreased seat belt use
Decreased helmet use
Reduced compliance with medication and treatment
Promiscuity
Hoarding
Choking sensation - globus
Real
Perceived
Respiratory/pulmonary
Asthma exacerbation
COPD
Increased risk of lung cancer
Hyperventilation/dyspnea
David McCollum, January 2009
ACADEMY ON VIOLENCE & ABUSE
ADVANCING HEALTH EDUCATION & RESEARCH
www.AVAHEALTH.org
384448_8x11.indd 1
7/31/09 11:08 AM